

Medications Containing Metformin

If you take any of these medications:

- **DO NOT** take it on the day of your CT scan
 - **DO NOT** take it for **48 hours after your scan**
- | | |
|---------------------------|---------------------------|
| • Actos Plus [®] | • Jentadueto [®] |
| • Avandamet [®] | • Kazano [®] |
| • Fortamet [®] | • Kombiglyze [®] |
| • Glucophage [®] | • Metaglip [®] |
| • Glucovance [®] | • Prandimet [®] |
| • Glumetza [®] | • Riomet [®] |
| • Invokamet [®] | • Synjardy [®] |
| • Janumet [®] | • Xigduo [®] |

The American College of Radiology recommends that your physician makes sure your kidneys are working normally before resuming your medication.

You received _____ of IV (intravenous) contrast today at Roswell Park for either a CT scan or an angiogram. Your kidneys will clear the contrast from your body, usually within 24 hours. In order to make sure your kidneys continue to work properly, it is important to drink extra fluids.

Oral Hydration

- **We recommend drinking at least 12 ounces (1.5 cups) of fluid every 2 hours for the next 24 hours.**
- Water is the best fluid to drink, but any caffeine-free beverage(s) is okay.
- Milk or any other dairy beverage is not recommended. (Dairy beverages are cleared from your body by your GI tract, not your kidneys.)

You may resume your medication on: _____